

Effect of the Theta-Beta Neurofeedback Protocol as a Function of Subtype in Children Diagnosed with Attention Deficit Hyperactivity Disorder

Elimelech Duarte Hernández, Javier González Marqués and Jesús M. Alvarado

Universidad Complutense (Spain)

Abstract. Neurofeedback is a neuronal self-regulation technique that teaches people to modulate their brain frequencies using visual and auditory reinforcements presented on a computer screen. To assess the effect of neurofeedback training in children with ADHD as far as improved attention and impulse control, and analyze whether or not there are differences between the inattentive and hyperactive subtypes. Fifty children diagnosed with ADHD participated in the study: 14 comprised the control group, and 36 the experimental group (16 with the inattentive ADHD subtype, 20 with the hyperactive ADHD subtype). Attention and impulse control were assessed using the Integrated Visual Auditory CPT (IVA/CPT). Results indicated that the predominantly inattentive group showed significant differences on the Control Scale ($p = .023$, $d = 1.31$) and the Attention Scale ($p < .01$, $d = 1.89$) of the IVA/CPT; meanwhile the predominantly hyperactive group showed significant improvement on the Control Scale ($p = .016$, $d = 1.21$). The control group exhibited no significant differences on either of the two scales ($p > .5$). In terms of theta/beta ratio, no significant differences were detected ($p = .10$) between ADHD subtypes. The findings suggest that neurofeedback training using the theta/beta protocol was more effective in the predominantly inattentive subset of individuals with ADHD.

Received 9 March 2015; Revised 30 April 2016; Accepted 2 May 2016

Keywords: ADHD subtypes, attention deficit disorder, EEG biofeedback, neurofeedback, theta/beta protocol.

The DSM-IV-TR describes Attention Deficit Hyperactivity Disorder (ADHD) as a persistent pattern of inattention and/or hyperactivity-impulsiveness that is more frequent or severe than individuals with a comparable level of development normally exhibit. According to the same diagnostic manual, the predominantly inattentive subtype meets the A1 criterion, but not A2, during the last six months; the predominantly hyperactive-impulsive subtype meets criterion A2, but not A1, during the last six months; and the combined subtype satisfies both criteria, A1 and A2, during the last six months (American Psychiatric Association, 2000). The current prevalence of Attention Deficit Disorder is estimated between 3 and 7% world-wide (Willcutt, 2012), and at 6.8% in Spain (Catalá-López et al., 2012).

Various studies have reported increased slow wave power in participants with ADHD compared to control groups, especially increased Theta (Amer, Rakhawy, & El Kholly, 2010; Chabot & Serfontein, 1996; Mann, Lubar, Zimmerman, Miller, & Muenchen, 1992) and Alpha band amplitude (Koehler et al., 2008; Lazzaro, Gordon, Li, Lim, & Plahn, 1999; Swartwood, Swartwood, Lubar, & Timmermann, 2003). Clarke and his colleagues have

additionally found a new encephalographic profile (EEG) relating to combined-subtype ADHD: a small increase in the proportion of Beta bands (Clarke, Barry, McCarthy, & Selikowitz, 2001b). Subsequent studies by the same research team at least support the existence of a specific EEG pattern in children with ADHD compared to a normative database (Barry, Clarke, McCarthy, & Selikowitz, 2002; Clarke et al., 2006), and differential EEG patterns in girls with predominantly inattentive and combined-type ADHD (Dupuy, Clarke, Barry, McCarthy, & Selikowitz, 2014). Meanwhile, Heinrich and his colleagues studied EEG activity in two groups of children with ADHD, one with the predominantly inattentive subtype and the other combined-type, while doing a task requiring attention. They discovered that the combined-type ADHD group had a high theta/alpha ratio, while the predominantly inattentive group had a high theta/beta ratio (Heinrich et al., 2014).

Neurofeedback, or EEG biofeedback, is a type of biofeedback that records electrical activity in the brain and transforms it into a digital visual and/or auditory signal, that is utilized as feedback, the purpose being to get the person to learn to self-regulate the amplitude of specific frequency waves (Hammond, 2011). The theta/beta neurofeedback training protocol has probably been the one most widely utilized in studies of neurofeedback and ADHD.

Correspondence concerning this article should be addressed to Elimelech Duarte Hernández. Instituto de Estudios Biofuncionales. Universidad Complutense. Paseo Juan XXIII, 11. 28040. Madrid (Spain).
E-mail: eduarte@neurotraining.es, javgonza@ucm.es

Some studies have reported significant improvements through neurofeedback training over controls (Drechsler et al., 2007; Gevensleben et al., 2009); improvement on tests of attention and response inhibition (Xiong, Shi, & Xu, 2005) and improvement in aspects of behavior and cognitive functioning (Nazari, Querne, De Broca, & Berquin, 2011).

Comparative studies have found that groups receiving neurofeedback treatment respond better than medicated groups in terms of inattentive behavior (Linden, Habib, & Radojevic, 1996; Monastrá, Monastrá, & George, 2002), yet others have reported significant improvements in symptomatology in neurofeedback as well as medicated groups (Fuchs, Birbaumer, Lutzenberger, Gruzelier, & Kaiser, 2003; Moreno-García, Delgado-Pardo, Camacho-Vara de Rey, Meneres-Sancho, & Servera-Barceló, 2015). On another note, Arns and his collaborators (Arns, de Ridder, Strehl, Breteler, & Coenen, 2009) completed a meta-analysis of 15 controlled studies ($N = 1194$), 11 of which used the theta/beta protocol. At the end of their analysis, they conclude that the neurofeedback technique is "efficacious and specific," and suggest that neurofeedback treatment of ADHD can be considered clinically significant, with a large effect size on symptoms of inattention and impulsivity, and a moderate effect size on hyperactivity symptoms. The same author later conducted a meta-analysis of the theta/beta ratio in cases of ADHD under the eyes open condition. This included nine studies ($N = 1253$) of children and adolescents 6 to 18 years old. From that analysis, the authors concluded that while a high theta/beta ratio cannot be considered a diagnostic measure of ADHD, it may have value in predicting how a patient will respond to pharmacological treatment versus neurofeedback treatment (Arns, Conners, & Kraemer, 2013). The protocol's main task is for subjects to learn to lower their theta band amplitude (8–12Hz) and increase their beta band amplitude (12–20Hz) (Lubar, 1997; Vernon et al., 2003). Using this protocol in clinical practice is supported by electrophysiology studies that have found significantly increased slow wave amplitude, and markedly decreased theta band amplitude in children with ADHD (Amer et al., 2010; Cornelio, Borbolla, & Gallegos, 2011; Helps et al., 2010; Yordanova, Heinrich, Kolev, & Rothenberger, 2006). Clarke, Barry, McCarthy, and Selikowitz (2001a) identified three different EEG groups in children with ADHD: the first characterized by excess slow wave activity and fast wave deficit, the second presenting significantly increased theta amplitude with decreased beta activity; and the third showed excess beta activity. The results led the authors to conclude that in terms of EEG profile, children with ADHD do not constitute a homogenous group (Clarke et al., 2001a).

The use of neurofeedback in clinical practice has increased in the past decade, bearing significant

improvement in some symptoms of ADHD (Butnik, 2005; Holtmann et al., 2004; Leins et al., 2007; Meisel, Servera, García-Banda, Cardo, & Moreno, 2013). Yet while the effectiveness of this neuronal self-regulation technique now has ample meta-analytical backing (Arns et al., 2009), few studies have examined differences between ADHD subtypes using a pre-post research design (Xiong et al., 2005), or compared the theta/beta protocol's effectiveness in different subtypes (Leins et al., 2007). To our understanding at the time this research was conducted, there was no evidence that any studies had investigated the theta/beta protocol's effects in ADHD subtypes by looking at EEG changes, or described the learning curve of training in cortical self-regulation of EEG waves.

The present study was conducted with the following objectives in mind: a) evaluate whether after neurofeedback training, differences are observed between a predominantly inattentive subgroup and a predominantly hyperactive ADHD subgroup in terms of IVA/CPT scores and theta/beta ratio; b) determine whether the theta-beta ratio behaves during training like a predictor variable of attention scores; and c) analyze whether the results obtained in the group that received neurofeedback training have some relationship with the form of medication used during training (immediate versus extended-release).

Method

Participants

Fifty-four children – 20 girls (average age 12.05; $SD = 2.49$) and 34 boys (average age 11.52; $SD = 3.05$) – diagnosed with ADHD participated in the study. Assignment to the neurofeedback treatment group was determined by participants' order of arrival. Personnel outside the research team prepared the participant list. Said personnel belonged to the center where the study was conducted, and the investigators were made aware of the participant list only at the end of the recruitment process. The first 40 participants were assigned to the neurofeedback (NF) treatment group ($n = 40$; average age = 11.53; $SD = 2.62$), of which 14 were girls (average age 11.81; $SD = 2.04$) and 26 were boys (average age = 11.38; $SD = 2.92$). The remaining 14 participants were assigned to the control group (average age = 12.25; $SD = 3.47$) – 6 girls (average age = 12.60; $SD = 3.50$) and 8 boys (average age = 11.98; $SD = 3.66$) – which received no intervention of any kind. Next, the group designated to receive neurofeedback training was divided into two subgroups according to their respective ADHD subtypes. In making that determination, we used the Spanish edition of the Behavior Assessment System for Children (BASC) (González-Marqués, Fernández-Guinea, Pérez-Hernández, & Santamaría, 2004) to identify

predominantly inattentive versus hyperactive cases. Combined-type cases were excluded from the study, as the investigators' main interest was to compare the effect of neurofeedback training on attentional variables and response inhibition. Thus, the composition of groups by subtype was as follows: the predominantly inattentive (PI) group ($n = 16$; average age = 11.57) was made up of 5 girls and 11 boys; and the predominantly hyperactive (PH) group ($n = 20$; average age = 11.91) was comprised of 6 girls and 14 boys. Of the participants in the treatment group, 14 were taking immediate-release (IR) methylphenidate (average age = 12.30), 24 were taking extended-release (ER) methylphenidate ($n = 24$; average age = 11.39), and two were taking different medications and were thus excluded from any analyses relating the form of drug to treatment results. All the children who participated belonged to the Association of Parents of Children with Attention Deficit with or without Hyperactivity of Madrid (ANSHDA from the acronym in Spanish). All participants were receiving pharmacological treatment at the time the study began and throughout neurofeedback training. To be accepted into the study, participants were required to meet all of the following inclusion criteria:

- Aged 7 to 17 years old
- Having a formal prior diagnosis of ADHD
- Keeping the medication dosage constant, and not changing it over the course of neurofeedback training.
- Intellectual quotient (IQ) > 80
- Not exhibiting other comorbid disorders
- Informed consent of parents

The parents of all children selected to participate in the study were informed in a private interview of their right to withdraw from participating in the study at any time. Written, signed consent forms were required from participants' parents before beginning treatment. The ADHD diagnosis and IQ of children chosen to participate in the study were confirmed by ANSHDA, where they were evaluated previously by competent professionals based on DSM-IV criteria. No additional assessments were applied for the purpose of diagnosis. Participants were randomly assigned to experimental groups.

This study was approved by the board of directors and the ethics committee of the ANSHDA., and was conducted in keeping with the Declaration of Helsinki (World Medical Association, 2013, October).

Materials and Procedure

Integrated Visual Auditory Continuous Performance Test (IVA/CPT)

The IVA/CPT is a computerized, standardized test developed to evaluate response inhibition and level of

attention (Sandford & Turner, 1994; Seckler, Burns, & Sandford, 1995, November). The test lasts approximately 13 minutes and mainly consists of 500 trials presenting visual and auditory patterns. Global quotient scores have a mean of 100 and a standard deviation of 15.

Scores on the six main scales of the IVA/CPT tap visual and auditory performance on two full scales: a) Full Scale Response Control Quotient (FSRCQ) scale, comprised of the subscales prudence, consistency, and stamina; and b) Full Scale Attention Quotient (FSAQ) scale, comprised of the subscales vigilance, focus, and speed. In the present study, we decided to utilize the global scales, demonstrated to be highly reliable measures (sensitivity 92%; specificity 90%; positive predictive strength 89%, and negative predictive strength 93%) of attention and impulse control (Sandford, Fine, & Goldman, 1995, Seckler et al., 1995) and validated measures that are helpful in diagnosing ADHD (Sandford & Turner, 1994). Furthermore, these global quotients provide stable measures over time, making them especially appropriate for studies with repeated measures (pre-post). Two IVA/CPT measures were taken from each participant: a) Before neurofeedback training started, and b) after neurofeedback training. The boys and girls who participated in the study were instructed to click on the button only when they saw or heard a "1," and not to click (inhibition) when they saw or heard a "2." IVA/CPT results are generally reported graphically as well as numerically so that changes in quotient scores can be calculated between the two times the IVA/CPT was administered and capture any treatment effects.

EEG frequency recording

Continuous data from pre-post EEG recordings were collected, and neurofeedback training carried out, using the BrainMaster System – Atlantis II. Specifically, the recording and training software utilized was BrainMaster 3.0 for clinical use, version 37i.

We took two EEG recordings (eyes open) for all children in a resting state, the first prior to neurofeedback training and the second 12.5 weeks later, following the last training session. In the control group, too, there was a period of 12.5 weeks between pre and post recordings. The sampling rate was 256 Hz, and the electrodes' impedance was kept below 5k Ω . A band-pass filter of 1 to 40 Hz was applied, and an electronic noise reduction filter, or *nocht* filter, was set at 50Hz. We utilized an exit channel on the central midline (CZ) a reference electrode placed frontally and centrally (FZ), and a ground electrode on the left earlobe (A1), according to the International 10–20 system. Cortical placement (CZ) was chosen for two reasons: first, previous studies have found significant results at that location associated

with neurofeedback training in cases of ADHD (Drechsler et al., 2007; Gevensleben et al., 2009; Heinrich, Gevensleben, Freisleder, Moll, & Rothenberger, 2004; Levesque, Beauregard, & Mensour, 2006). Furthermore, we believe CZ is appropriate given that its placement is relatively free from eye movements and EMG artifacts compared to other sites closer to the eyes and jawbones; it is beneficial to keep that in mind when training children with ADHD. Electrooculography (EOG) and electromyography (EMG) were not used, because the EEG recording software has integrated filters (120 μ V) that detect and exclude signals over the threshold, excluding EOG and EMG artifacts. We also visually inspected the data prior to analysis to confirm the absence of artifacts. Pre- and post-training EEG recording data were obtained from global EEG recordings during neurofeedback training. Individual averages were calculated from the theta and beta band amplitudes recorded during each session (25 recordings of 60 seconds each). Averages for all participants were subsequently sorted.

Neurofeedback training protocol

Neurofeedback training was carried out in an isolated room, without interruption. All the children who participated in the study received two weekly neurofeedback sessions of 25 minutes each for 12.5 weeks, completing a total of 25 sessions. The neurofeedback training protocol was designed to reduce Theta band activity and increase Beta band activity.

The neurofeedback training interface was presented to participants on a computer screen in the form of eight different types of digital games, synchronized with a preset training protocol. The games were presented progressively over the course of the 25 training sessions so as to maintain participants' motivation for the full duration of training. All the games had a visual display showing the respondent's score, updated in real time, in the upper left or right of the screen. It was synchronized with an auditory stimulus they heard through the speaker whenever they met the task conditions simultaneously: a) to lower theta-band amplitude, and b) increase beta-band amplitude. A reward criterion was set: one point for every 500 milliseconds that the task lasted.

Statistical analyses

To statistically analyze all the measures taken, the SPSS statistical package, version 20, was utilized. To evaluate the effect of training on different IVA/CPT measures, average scores in the experimental and control groups were compared using Student's *t* test for repeated measures. This first study was completed using regression analysis, the purpose being to test the predictive hypothesis that children who receive neurofeedback

training will score higher on the IVA/CPT as a function of having achieved better neuronal self-regulation in terms of theta/beta ratio. In cases where Student's *t* indicated statistically significant differences ($p < .05$) between pre- and post-treatment means, effect size was computed per Cohen: $d = M / SD$. We took into account the following interpretation criteria: small: $|d| = .20$ to $.50$; medium: $|d| = .50$ to $.80$; and large: $|d| \geq .80$ (Cohen, 1998). To avoid overestimating effect size, we used the Standard Deviation (*SD*) of the test rather than *SD* of the differences between pre and post measures (Dunlap, Cortina, Vaslow, & Burke, 1996). Finally, repeated measures Analysis of Variance (ANOVA) was used to test whether factors like ADHD subtype, medication type, or their interaction, were related to success in neurofeedback training.

Results

Pre-post IVA/CPT Measurement in the Control Group vs. Neurofeedback Group

As shown in Table 1, the means difference in the control group was negative, because post IVA/CPT scores were slightly lower than pre; in any case, the differences were not statistically significant on the FSAQ ($M = -3.50$, $SD = 13.6$), $t(13) = -.97$, $p = .352$, $d = -0.535$ nor the FSRCQ ($M = -0.86$, $SD = 28.2$), $t(13) = -.11$, $p = .911$, $d = -0.063$. That is consistent with previous studies' results in terms of IVA/CPT measures, which show good temporal stability and pre-post reliability in the absence of some mediating intervention (Sandford & Turner, 1994).

On another note, with respect to the groups who received neurofeedback training, significant improvements were observed on the FSRCQ in the predominantly inattentive group ($M = 8.12$, $SD = 13.2$), $t(15) = 2.54$, $p = .023$, $d = 1.31$, as well as the predominantly hyperactive group ($M = 5.90$, $SD = 9.8$), $t(19) = 2.63$, $p = .016$, $d = 1.21$ (see Table 1). Referring to the FSAQ, significant differences were observed in the predominantly inattentive group ($M = 19.35$, $SD = 21.8$), $t(15) = 3.66$, $p < .01$, $d = 1.89$, whereas differences in the predominantly hyperactive group did not reach the level of statistical significance ($M = 7.105$, $SD = 18.1$), $t(19) = 1.71$, $p > .04$, $d = 0.79$.

Efficacy of the Intervention as a Function of Subtype and Form of Medication

To evaluate possible effects of the variables ADHD subtype and type of medication on treatment efficacy, repeated measures ANOVA was carried out. The measurement factor was the 25 sessions averaged across five periods: Period 1, sessions 1 to 5; Period 2, sessions 6 to 10; Period 3, sessions 11 to 15; Period 4, sessions 16 to 20, and Period 5, sessions 21 to 25. The independent

Table 1. *T Test for Related Samples in the Control Group and Groups that Received Neurofeedback Training*

| Group | Variable | Related Differences | | | |
|---------------------|---------------------|---------------------|------|------------|--------|
| | | Mean | SD | Standard E | t |
| Control | Response Inhibition | -0.86 | 28.2 | 7.53 | -0.11 |
| | Attention | -3.50 | 13.6 | 3.63 | -0.97 |
| Hyperactive Subtype | Response Inhibition | 5.90 | 9.8 | 2.24 | 2.63* |
| | Attention | 7.11 | 18.1 | 4.15 | 1.71 |
| Inattentive Subtype | Response Inhibition | 8.12 | 13.2 | 3.19 | 2.54* |
| | Attention | 19.35 | 21.8 | 5.29 | 3.66** |

**p* < .05.

***p* < .01.

variables were: a) ADHD subtype – Inattentive Subtype (IS) or Hyperactive Subtype (HS); and b) Form of methylphenidate – Immediate Release (IR) or Extended Release (ER). Additionally, a post-hoc pairwise comparison analysis was done, based on Student's *t* test and implemented in SPSS – called the Minimum Significant Difference (MSD) – in order to determine statistical significance between levels of the factors.

ANOVA revealed a significant main effect of the treatment variable, $F(4, 128) = 6.578, p < .001$, partial $\eta^2 = .17$, and an interaction effect of treatment and ADHD subtype, $F(4, 128) = 3.536, p = .009$, partial $\eta^2 = .10$. Statistically significant differences were not observed according to ADHD subtype, $F(1, 32) = 2.869, p = .10$, or type of medication, $F(1, 34) = 1.470, p = .234$.

Training produced a gradual decrease in average theta/beta ratio in the five periods analyzed, but the most marked decrease occurred during the first three periods (the first 15 sessions): means $S1_5 = 2.18, S6_10 = 2.04, S11_15 = 1.81, S16_20 = 1.78, S21_25 = 1.77$. Post-hoc pairwise comparison through MSD confirmed that between the first two periods and the rest, differences were generally statistically significant (see Table 2).

Figure 1 represents the interaction between neurofeedback training and ADHD subtype. It shows a steep decrease in theta/beta ratio with training during the first three periods (first 15 sessions) in the predominantly inattentive group. Conversely, the predominantly hyperactive group exhibited a slower, more gradual decrease all the way through the final period considered (sessions 21 to 25).

Regarding medication type, no significant differences were found in theta/beta ratio between subjects taking Immediate Release (IR) versus Extended Release (ER) methylphenidate.

Relationship between Lower Theta/beta Ratio and Improved Attention

Table 3 presents results from the regression model, using average theta/beta ratios from the first five and last five

Table 2. *Pairwise Comparison of Theta/Beta Ratio in the Five Training Periods Analyzed*

| Period 1 | Period 2 | Difference (1-2) | Stand. E. | <i>p</i> |
|----------|----------|------------------|-----------|----------|
| S1_5 | S6_10 | 0.143 | 0.085 | .104 |
| | S11_15 | 0.366 | 0.128 | .007* |
| | S16_20 | 0.401 | 0.140 | .007* |
| | S21_25 | 0.414 | 0.134 | .004* |
| S6_10 | S11_15 | 0.224 | 0.109 | .049* |
| | S16_20 | 0.258 | 0.123 | .043* |
| | S21_25 | 0.271 | 0.115 | .025* |
| S11_15 | S16_20 | 0.034 | 0.026 | .198 |
| | S21_25 | 0.047 | 0.024 | .054 |
| S16_20 | S21_25 | 0.013 | 0.026 | .068 |

Note: The measurement factor was the 25 sessions averaged across five periods: Period 1: S1_5; Period 2: S6_10; Period 3: S11_15; Period 4: S16_20, and Period 5: S21_25.

**p* < .05.

***p* < .01.

sessions as predictor variables. Results indicate that these variables predict attention scores in the form of a linear model with predictive validity $r = .54$.

$$\text{Attention} = 128.88 + 14.17 (S1_5) - 40.18 (S21_25)$$

The scatter plot in Figure 2 conveys that values on the FSAQ increase as a function of the difference between the first period S1_5 and last period S20_25.

Discussion

After carrying out a training program using the neurofeedback technique and applying the theta/beta protocol to two groups of children diagnosed with ADHD, one predominantly inattentive and one predominantly hyperactive, our results verify that the predominantly inattentive group obtained better results




Figure 1. Interaction Effect of Neurofeedback Training and ADHD Subtype.

Note: Notice that in the figure, the inattentive group’s theta/beta ratio decreases more steeply than in the hyperactive group, which shows a more gradual curve.

Table 3. Regression Model of Theta/Beta Ratio’s Ability to Predict Attentional Performance

| Coefficients ^a | | | | | |
|---------------------------|-----------------------------|----------------|---------------------------|-------|--------|
| Sessions | Unstandardized Coefficients | | Standardized Coefficients | | |
| | B | Standard Error | β | t | p |
| (Constant) | 128.88 | 14.54 | | 8.87 | < .001 |
| S1 – S5 | 14.17 | 5.62 | .487 | 2.52 | .017 |
| S21 – S25 | -40.18 | 10.82 | -.717 | -3.71 | .001 |

^adependent variable: performance on FSAQ.

than the predominantly hyperactive group. On the other hand, our analyses suggest that theta/beta ratio behaves like variable that predicts scores on attention scales.

In this study’s first objective, we sought to determine whether after theta-beta protocol training, differences would be observed in measures of attentional variables between a group of predominantly hyperactive children, and a group of predominantly inattentive children.

The main purpose of that objective relates to IVA/CPT results, in particular, to identify potential variations between pre- and post-treatment measures on the test’s main scales. Regarding the FSRCQ, we observed significant differences in the predominantly inattentive group ($p < .05$) as well as the predominantly hyperactive group ($p < .05$), the result being particularly favorable in the former group since the FSRCQ relates to inhibitory control, consistency, and stamina. Those results are consistent with Xiong’s findings, who utilized the same test and found improvement on the FSRCQ for both subtypes, following a neurofeedback training phase (Xiong et al., 2005). Other studies that also utilized the IVA/CPT before and after neurofeedback training, but did not discriminate among ADHD subtypes, have reported improvements on both scales (Moreno-García et al., 2015), or on FSAQ (Hillard, El-Baz, Sears, Tasman, & Sokhadze, 2013).

A notable result regarding the FSAQ is that only the predominantly inattentive group showed significant differences between pre- and post-treatment measures ($p = .02$), and the effect size was large ($d = 1.88$). That is not in line with Xiong’s results, who found improvement in groups made up of both subtypes. The discrepancy between our results and Xiong’s may relate to the number




Figure 2. Scatter Plot of the Regression Model with Predictor Variable in Attention.

Note: The values in the attention scale increase as a function of the difference between the first period S1_S5 (sessions 1 to 5) and last period S20_25 (sessions 20 to 25).

of neurofeedback sessions used in that study compared to ours. If that relationship is confirmed through controlled studies, including a larger number of sessions, it would support the hypothesis that it takes more time for children with predominantly hyperactive ADHD to learn to self-regulate attention-related neuronal response than to improve self-regulation of response inhibition. The analysis of variance we conducted – to determine if the efficacy of neurofeedback training was related to ADHD subtype – seems to back that hypothesis. Taking theta/beta ratio as a variable indicative of training, we performed repeated measures analysis of variance, comparing five periods, each corresponding to a five-session phase of total training. Results suggested that in the predominantly inattentive group of children, theta/beta ratio declined steeply during the first 15 sessions, then more gradually until the 25 sessions were over. Conversely, the predominantly hyperactive group displayed a subtler decrease in theta/beta ratio across the entire course of training, with steady slope until the final session. This suggests that predominantly hyperactive children need a greater number of sessions to lower their theta/beta ratio, at least in the case of the protocol studied here. These results are not conclusive, but they do indicate it is important to more closely examine the behavior of the theta/beta protocol

learning curve to corroborate this finding in future studies with a greater number of sessions.

The second objective of this study was to evaluate whether there is a correlation between IVA/CPT results in the group receiving neurofeedback training, and their level of neuronal self-regulation of theta/beta ratio during the 25 training sessions. That analysis' point of interest lies in estimating whether theta/beta ratio behaves like a predictor variable of scores on the FSRCQ and FSAQ scales of the IVA/CPT. Results obtained through regression analysis indicate that theta/beta ratio indeed predicts attention scores, fitting a linear model, with predictive validity $r = .54$. These results could be of clinical utility in cases where cerebral mapping prior to neurofeedback training is not available, because with such a map, the EEG profile of a child with ADHD can be identified to see if it fits the profile of excess theta activity (Chabot & Serfontein, 1996) and deficient beta frequency, which is the profile predominantly found in cases of ADHD (Amer et al., 2010; Mann et al., 1992), profiles where beta waves predominate (Clarke et al., 2001b), or profiles with irregularities in other brain waves (Koehler et al., 2008; Lazzaro et al., 1999; Swartwood et al., 2003). However, in cases where the theta/beta protocol is applied and there is no previous electroencephalographic test to go on,

one can use theta/beta curve analysis after an initial block of 15 to 20 sessions and estimate whether or not the protocol will be effective in this specific case.

Finally, regarding the third objective, which sought to determine if there was a relationship between the neurofeedback training group's results and the form of pharmacological treatment, significant differences were not observed between participants taking the Immediate Release (IR) versus Extended Release (ER) form.

This study contributes to our understanding of how neurofeedback training using the theta/beta protocol influences response inhibition and attentional variables in cases of ADHD, with either inattentive or hyperactive predominance. Furthermore, it makes a new contribution to the clinical field in suggesting the theta/beta ratio can behave like a predictor variable of scores on attention scales. Nevertheless, this study's limitations, such as sample size, sampling method, and the wide age range of participants, mean these results must be interpreted with caution and moderation. That being said, the results reported here serve as a foundation for conducting future research geared toward optimizing the use of neurofeedback intervention protocols.

References

- Amer D. A., Rakhawy M. Y., & El Kholly S. H. (2010). Quantitative EEG in children with Attention Deficit Hyperactivity Disorder. *The Egyptian Journal of Neurology and Neurosurgery*, 47, 399–406.
- American Psychiatric Association (2000). *Diagnostic and statistical manual of mental disorders* (4th Ed., Text Rev.). Washington, DC: Author.
- Arns M., Conners C. K., & Kraemer H. C. (2013). A decade of EEG Theta/Beta ratio research in ADHD: A meta-analysis. *Journal of Attention Disorders*, 17, 374–383. <http://dx.doi.org/10.1177/1087054712460087>
- Arns M., de Ridder S., Strehl U., Breteler M., & Coenen A. (2009). Efficacy of neurofeedback treatment in ADHD: The effects on inattention, impulsivity and hyperactivity: A meta-analysis. *Clinical EEG and Neuroscience*, 40, 180–189. <http://dx.doi.org/10.1177/155005940904000311>
- Barry R. J., Clarke A. R., McCarthy R., & Selikowitz M. (2002). EEG coherence in attention-deficit/hyperactivity disorder: A comparative study of two DSM-IV types. *Clinical Neurophysiology*, 113, 579–585. [http://dx.doi.org/10.1016/S1388-2457\(02\)00036-6](http://dx.doi.org/10.1016/S1388-2457(02)00036-6)
- Butnik S. M. (2005). Neurofeedback in adolescents and adults with attention deficit hyperactivity disorder. *Journal of Clinical Psychology*, 61, 621–625. <http://dx.doi.org/10.1002/jclp.20124>
- Catalá-López F., Peiró S., Rídao M., Sanfélix-Gimeno G., Gènova-Maleras R., & Catalá M. A. (2012). Prevalence of attention deficit hyperactivity disorder among children and adolescents in Spain: A systematic review and meta-analysis of epidemiological studies. *BMC Psychiatry*, 12, 168. <http://dx.doi.org/10.1186/1471-244x-12-168>
- Clarke A. R., Barry R. J., McCarthy R., & Selikowitz M. (2001a). EEG-defined subtypes of children with attention-deficit/hyperactivity disorder. *Clinical Neurophysiology*, 112, 2098–2105. [http://dx.doi.org/10.1016/S1388-2457\(01\)00668-X](http://dx.doi.org/10.1016/S1388-2457(01)00668-X)
- Clarke A. R., Barry R. J., McCarthy R., & Selikowitz M. (2001b). Excess beta activity in children with attention-deficit/hyperactivity disorder: An atypical electrophysiological group. *Psychiatry Research*, 103, 205–218. [http://dx.doi.org/10.1016/S0165-1781\(01\)00277-3](http://dx.doi.org/10.1016/S0165-1781(01)00277-3)
- Clarke A. R., Barry R. J., McCarthy R., Selikowitz M., Magee C. A., Johnstone S. J., & Croft R. J. (2006). Quantitative EEG in low-IQ children with attention-deficit/hyperactivity disorder. *Clinical Neurophysiology*, 117, 1708–1714. <http://dx.doi.org/10.1016/j.clinph.2006.04.015>
- Cohen J. D. (1998). *Statistical power analysis for the behavioral sciences* (2nd Ed.). Hillsdale, NY: Erlbaum.
- Cornelio-Nieto J., Borbolla-Sala M. E., & Gallegos-Dimas A. (2011). Alteraciones electroencefalográficas en niños con trastorno por déficit de atención/hiperactividad [Electroencephalographic alterations in children with attention deficit hyperactivity disorder]. *Revista de Neurología*, 52, S97–S101.
- Chabot R. J., & Serfontein G. (1996). Quantitative electroencephalographic profiles of children with attention deficit disorder. *Biological Psychiatry*, 40, 951–963. [http://dx.doi.org/10.1016/0006-3223\(95\)00576-5](http://dx.doi.org/10.1016/0006-3223(95)00576-5)
- Drechsler R., Straub M., Doehner M., Heinrich H., Steinhausen H. C., & Brandeis D. (2007). Controlled evaluation of a neurofeedback training of slow cortical potentials in children with Attention Deficit/Hyperactivity Disorder (ADHD). *Behavioral and Brain Functions*, 3, 35. <http://dx.doi.org/10.1186/1744-9081-3-35>
- Dunlap W. P., Cortina J. M., Vaslow J. B., & Burke M. J. (1996). Meta-analysis of experiments with matched groups or repeated measures designs. *Psychological Methods*, 1, 170–177. <http://dx.doi.org/10.1037/1082-989X.1.2.170>
- Dupuy F. E., Clarke A. R., Barry R. J., McCarthy R., & Selikowitz M. (2014). EEG differences between the combined and inattentive types of attention-deficit/hyperactivity disorder in girls: A further investigation. *Clinical EEG and Neuroscience*, 45, 231–237. <http://dx.doi.org/10.1177/1550059413501162>
- Fuchs T., Birbaumer N., Lutzenberger W., Gruzelier J. H., & Kaiser J. (2003). Neurofeedback treatment for attention deficit hyperactivity disorder in children: A comparison with methylphenidate. *Applied Psychophysiology and Biofeedback*, 28, 1–12.
- Gevensleben H., Holl B., Albrecht B., Schlamp D., Kratz O., Studer P., ... Heinrich H. (2009). Distinct EEG effects related to neurofeedback training in children with ADHD: A randomized controlled trial. *International Journal of Psychophysiology*, 74, 149–157. <http://dx.doi.org/10.1016/j.jpsycho.2009.08.005>
- González-Marqués J., Fernández-Guinea S., Pérez-Hernández E., & Santamaría P. (2004). *BASC. Sistema de Evaluación de la Conducta de Niños y Adolescentes* [BASC. Behavior Assessment System for Children and Adolescents]. Madrid, Spain: TEA Ediciones.

- Hammond D. C.** (2011). What is neurofeedback: An update. *Journal of Neurotherapy: Investigations in Neuromodulation, Neurofeedback and Applied Neuroscience*, 15, 305–336. <http://dx.doi.org/10.1080/10874208.2011.623090>
- Heinrich H., Busch K., Studer P., Erbe K., Moll G. H., & Kratz O.** (2014). EEG spectral analysis of attention in ADHD: Implications for neurofeedback training? *Frontiers in Human Neuroscience*, 8, 611. <http://dx.doi.org/10.3389/fnhum.2014.00611>
- Heinrich H., Gevensleben H., Freisleder F. J., Moll G. H., & Rothenberger A.** (2004). Training of slow cortical potentials in attention-deficit/hyperactivity disorder: Evidence for positive behavioral and neurophysiological effects. *Biological Psychiatry*, 55, 772–775. <http://dx.doi.org/10.1016/j.biopsych.2003.11.013>
- Helps S. K., Broyd S. J., James C. J., Karl A., Chen W., & Sonuga-Barke E. J. S.** (2010). Altered spontaneous low frequency brain activity in Attention Deficit/Hyperactivity Disorder. *Brain Research*, 1322, 134–143. <http://dx.doi.org/10.1016/j.brainres.2010.01.057>
- Hillard B., El-Baz A. S., Sears L., Tasman A., & Sokhadze E. M.** (2013). Neurofeedback training aimed to improve focused attention and alertness in children with ADHD: A study of relative power of EEG rhythms using custom-made software application. *Clinical EEG and Neuroscience*, 44, 193–202. <http://dx.doi.org/10.1177/1550059412458262>
- Holtmann M., Stadler C., Leins U., Strehl U., Birbaumer N., & Poustka F.** (2004). Neurofeedback for the treatment of attention-deficit/hyperactivity disorder (ADHD) in childhood and adolescence. *Zeitschrift für Kinder - und Jugendpsychiatrie und Psychotherapie*, 32, 187–200. <http://dx.doi.org/10.1024/1422-4917.32.3.187>
- Koehler S., Lauer P., Schreppe T., Jacob C., Heine M., Boreatti-Hümmer A., ... Herrmann M. J.** (2009). Increased EEG power density in alpha and theta bands in adult ADHD patients. *Journal of Neural Transmission*, 116(1), 97–104. <http://dx.doi.org/10.1007/s00702-008-0157-x>
- Lazzaro I., Gordon E., Li W., Lim C. L., Plahn M., Whitmont S., ... Meares R.** (1999). Simultaneous EEG and EDA measures in adolescent Attention Deficit Hyperactivity Disorder. *International Journal of Psychophysiology*, 34, 123–134. [http://dx.doi.org/10.1016/S0167-8760\(99\)00068-9](http://dx.doi.org/10.1016/S0167-8760(99)00068-9)
- Leins U., Goth G., Hinterberger T., Klinger C., Rumpf N., & Strehl U.** (2007). Neurofeedback for children with ADHD: A comparison of SCP and Theta/Beta protocols. *Applied Psychophysiology and Biofeedback*, 32, 73–88. <http://dx.doi.org/10.1007/s10484-007-9031-0>
- Levesque J., Beauregard M., & Mensour B.** (2006). Effect of neurofeedback training on the neural substrates of selective attention in children with attention-deficit/hyperactivity disorder: A functional magnetic resonance imaging study. *Neuroscience Letters*, 394, 216–221. <http://dx.doi.org/10.1016/j.neulet.2005.10.100>
- Linden M., Habib T., & Radojevic V.** (1996). A controlled study of the effects of EEG biofeedback on cognition and behavior of children with attention deficit disorder and learning disabilities. *Biofeedback and Self-Regulation*, 21, 297. <http://www.dx.doi.org/10.1007/BF02214740>
- Lubar J. F.** (1997). Neocortical dynamics: Implications for understanding the role of neurofeedback and related techniques for the enhancement of attention. *Applied Psychophysiology and Biofeedback*, 22, 111–126. <http://dx.doi.org/10.1023/A:1026276228832>
- Mann C. A., Lubar J. F., Zimmerman A. W., Miller C. A., & Muenchen R. A.** (1992). Quantitative analysis of EEG in boys with attention-deficit-hyperactivity disorder: Controlled study with clinical implications. *Pediatric Neurology*, 8(1), 30–36. [http://dx.doi.org/10.1016/0887-8994\(92\)90049-5](http://dx.doi.org/10.1016/0887-8994(92)90049-5)
- Meisel V., Servera M., Garcia-Banda G., Cardo E., & Moreno I.** (2013). Neurofeedback and standard pharmacological intervention in ADHD: A randomized controlled trial with six-month follow-up. *Biological Psychology*, 94(1), 12–21. <http://dx.doi.org/10.1016/j.biopsycho.2013.04.015>
- Monastra V. J., Monastra D. M., & George S.** (2002). The effects of stimulant therapy, EEG biofeedback, and parenting style on the primary symptoms of Attention-Deficit/Hyperactivity Disorder. *Applied Psychophysiology and Biofeedback*, 27, 231–249. <http://dx.doi.org/10.1023/A:1021018700609>
- Moreno-García I., Delgado-Pardo G., Camacho-Vara de Rey C., Meneres-Sancho S., & Servera-Barceló M.** (2015). Neurofeedback, pharmacological treatment and behavioral therapy in hyperactivity: Multilevel analysis of treatment effects on electroencephalography. *International Journal of Clinical and Health Psychology*, 15, 217–225. <http://dx.doi.org/10.1016/j.ijchp.2015.04.003>
- Nazari A. M., Querne L., De Broca A., & Berquin P.** (2011). Effectiveness of EEG biofeedback as compared with methylphenidate in the treatment of Attention-Deficit/Hyperactivity Disorder: A clinical outcome study. *Neuroscience & Medicine*, 2, 78–86. <http://dx.doi.org/10.4236/nm.2011.22012>
- Sandford J. A., Fine A. H., & Goldman L.** (1995, August). Validity study of the IVA: A visual and auditory CPT. *Paper presented at the annual convention of the American Psychological Association*, New York, NY.
- Sandford J. A., & Turner A.** (1994). *Manual for the Integrated Visual and Auditory Continuous Performance Test V*. Richmond, VA: Brain Train.
- Seckler P., Burns W., & Sandford J. A.** (1995, November). A reliability study of IVA: Intermediate Visual and Auditory Continuous Performance Test. *Paper presented at the 1995 annual convention of CHADD*, Washington, DC.
- Swartwood J. N., Swartwood M. O., Lubar J. F., & Timmermann D. L.** (2003). EEG differences in ADHD-combined type during baseline and cognitive tasks. *Pediatric Neurology*, 28, 199–204. [http://dx.doi.org/10.1016/S0887-8994\(02\)00514-3](http://dx.doi.org/10.1016/S0887-8994(02)00514-3)
- Vernon D., Egner T., Cooper N., Compton T., Neilands C., Sheri A., & Gruzelier J.** (2003). The effect of training distinct neurofeedback protocols on aspects of cognitive performance. *International Journal of Psychophysiology*, 47(1), 75–85. [http://dx.doi.org/10.1016/S0167-8760\(02\)00091-0](http://dx.doi.org/10.1016/S0167-8760(02)00091-0)
- Willcutt E. G.** (2012). The prevalence of DSM-IV attention-deficit/hyperactivity disorder: A meta-analytic review. *Neurotherapeutics*, 9, 490–499. <http://dx.doi.org/10.1007/s13311-012-0135-8>

World Medical Association (2013, October). WMA Declaration of Helsinki - ethical principles for medical research involving human subjects. *Paper presented at the 64th WMA General Assembly*, Fortaleza, Brazil.

Xiong Z., Shi S., & Xu H. (2005). A controlled study of the effectiveness of EEG biofeedback training on children with attention deficit hyperactivity disorder. *Journal of Huazhong University of Science and*

Technology Medical Sciences, 25, 368–370. <http://dx.doi.org/10.1007/BF02828171>

Yordanova J., Heinrich H., Kolev V., & Rothenberger A. (2006). Increased event-related theta activity as a psychophysiological marker of comorbidity in children with tics and Attention-Deficit/Hyperactivity Disorders. *NeuroImage*, 32, 940–955. <http://dx.doi.org/10.1016/j.neuroimage.2006.03.056>